

Electroweak Precision Measurements in Diboson Production at CMS

Pietro Vischia

(on behalf of the CMS Collaboration)

*Centre for Cosmology, Particle Physics and Phenomenology - CP3, Université catholique de Louvain, 2
Chemin du Cyclotron - Box L7.01.05, B-1348 Louvain-la-Neuve, Belgium*




In this contribution, I have outlined recent precision measurements of the standard model (SM) multiboson production at CMS. A study of diboson production at 5 TeV constitutes an important probe of the SM at a new energy, and the data favour NNLO predictions obtained by MATRIX. A study of WZ production at 13 TeV constitutes the most comprehensive study of WZ production to date, containing inclusive and differential cross section measurements, charge asymmetry measurements, constraints on the LHC proton parton distribution functions, and constraints on anomalous values of the WWZ trilinear gauge coupling. No evidence for new physics is found, and all the results favour SM predictions calculated at NNLO using MATRIX.

1 Multiboson Production at CMS

The associated production of two or three vector bosons (W or Z) at the LHC constitutes a set of important processes that help shed light on the standard model (SM) of particle physics. On one side, multiboson production constitutes an important background to measurements of the Higgs boson properties or to searches of new physics. On the other side new physics may appear directly in multiboson production in the form of anomalous values of the triple and quadruple gauge bosons couplings that intervene in the production of these processes. In this Manuscript, I will describe the most recent precision measurement of SM multiboson production performed by the CMS Collaboration with LHC data, namely the study of diboson production at a centre-of-mass energy of 5 TeV¹ and the study of electroweak (EWK) WZ production at 13 TeV². Associated WZ production via vector boson scattering is the focus of the contributions by Oleg Kuprash and Mattia Lizzo at this very same conference.

2 Diboson production at 5 TeV

Diboson (WW, WZ, and ZZ) production constitutes an important probe for the dependence of SM cross sections on the beam energy; measuring diboson production cross sections at a center-of-mass energy of 5 TeV is therefore of paramount importance. The CMS analysis¹ makes use of



Process	Estimation	Total cross section [pb]
WW	MATRIX	$29.8^{+0.7}_{-0.6}$ (scale)
	Measured	$37.0^{+5.5}_{-5.2}$ (stat) $^{+2.7}_{-2.6}$ (syst)
WZ	MATRIX	$11.3^{+0.2}_{-0.2}$ (scale)
	Measured	$6.4^{+2.5}_{-2.1}$ (stat) $^{+0.5}_{-0.3}$ (syst)
ZZ	MATRIX	$3.9^{+0.1}_{-0.1}$ (scale)
	Measured	$5.3^{+2.5}_{-2.1}$ (stat) $^{+0.5}_{-0.4}$ (syst)

Figure 1 – Evolution of VV cross section measurements as a function of the centre-of-mass energy (left). Measured cross sections for VV production, compared with predictions from MATRIX³ (right). Figure and table reproduced from the CMS paper¹.




Figure 2 – Feynman diagrams for WZ production. The diagram on the left is sensitive to anomalous values of the WWZ coupling. Figures reproduced from the CMS paper².

single-lepton triggers to study diboson production in a final state characterized by the presence of multiple leptons: WW production is studied by requiring two opposite-charge different-flavour leptons, plus transverse mass requirements and vetoing jet production; WZ production is targeted in a three-lepton and in a two-muons same-sign region with additional requirements that identify the pairs of leptons that most likely originate from the decay of a Z boson; and the ZZ is studied in a four-lepton region and in a two-lepton opposite-charge same-flavour requiring the missing transverse energy p_T^{miss} to be $p_T^{\text{miss}} > 50\text{GeV}$. All the background contributions are estimated from simulation, except for the contribution from objects misidentified as leptons (*nonprompt lepton* contribution), which is estimated from data. The results, illustrated and tabulated in Fig. 1, are well in agreement with NNLO predictions from MATRIX³, except in WZ, where the agreement is within two standard deviations.

3 WZ production at 13 TeV

The associated production of a W and a Z boson at 13 TeV has been studied using the full CMS Run II data set². This process features a charged final state that is sensitive to the quark parton distribution functions (PDFs)⁴, because at the tree level it is completely dominated by $q\bar{q}'$ states. WZ is also sensitive at tree level to anomalous values of the WWZ triple gauge coupling, and it is the dominant SM background process in any analysis targeting trilepton final states with low hadronic activity, such as $t\bar{t}H$ ⁵, $t\bar{t}V$ (the first full Run 2 determination of $t\bar{t}W$ cross section has been published in⁵), supersymmetric electroweak searches⁶, etc. The diagrams for WZ production are shown in Fig. 2.

This CMS study² is a comprehensive study of WZ production, leveraging the multilepton final state to measure fiducial, inclusive, and differential cross sections, vector boson polarization,

and the value of the trilinear gauge coupling that intervenes in Fig. 2 (left). For the first time, WZ production is used to probe CP-violating parameters in effective field theory (EFT) rather than the traditional CP-conserving ones, and we use Bayesian methods to innovatively constrain the proton PDFs predicted by the LHAPDF Collaboration⁴.

The analysis uses a set of single-lepton and dilepton triggers with an overall efficiency of about 100%, and with respect to previous publications⁷, we have improved the lepton reconstruction to provide additional background suppression. All leptons are *dressed*, i.e. the photon momenta in a radius of 0.10 in the (η, ϕ) plane are added to the lepton momentum, and the dedicated boosted decision tree that identifies leptons has been now retuned based on optimizations we performed in the context of the CMS Observation of ttH production⁸. For electrons, a tight-charge criterium is also included: it reduces the acceptance by $< 1\%$ while reducing the background contribution from *charge flips*, i.e. misreconstructed electron charge, from 0.2% to 0.03%. We compute lepton identification efficiencies⁹ in simulated events, using WZ production as a signal, and the set of all nonprompt background contributions—mostly $t\bar{t}$ and Drell-Yan—as background, according to the formula:

$$\epsilon(MVA) := \frac{\#\text{events in SR(Loose + MVA cut)}}{\#\text{events in SR(loose)}}. \quad (1)$$

The event selection consists of a baseline selection of three light leptons (electrons or muons), out of which an opposite-sign same-flavour pair compatible with the hypothesis of coming from a Z boson decay is required. A no-heavy-flavour-activity, high-trilepton-mass signal region (SR) is used in conjunction with three control regions (CRs) where we invert some of the signal region requirements to target specific backgrounds: a ZZ CR is characterized by requiring four leptons, a Conversions CR by requiring low p_T^{miss} and dilepton mass, and a $t\bar{t}$ CR by requiring the presence of jets coming from the fragmentation of b quarks and no Z mass window requirement. We compute the fiducial cross sections by means of a maximum likelihood fit to the lepton flavour composition in each of the regions while also constraining the normalization of the three backgrounds in their CRs. For the ZZ region, the lepton flavour is taken as the flavour of the three leading- p_T leptons, to mimic what would affect us in the SR if we didn't reconstruct the fourth lepton.

We find that the observed data favour the NNLO predictions from MATRIX³, and we extrapolate the fiducial results to the inclusive phase space accounting for leptonic branching fractions. Our results, displayed in Fig. 3 and tabulated in the paper², are affected by a 4% overall uncertainty on each lepton flavour final state, which is smaller than the best determination to date (5% by the ATLAS Collaboration¹⁰). The dominant systematic uncertainty components are, in decreasing order, the beam luminosity, the b quark tagging, and the nonprompt lepton identification, whereas the statistical uncertainty accounts for about 90% of the total uncertainty. Combining the result across the four lepton flavour final states yields a cross section measured with an overall uncertainty of 3%, which is about half of similar measurements by ATLAS¹⁰ and CMS⁷ and is below the theoretical uncertainties in the POWHEG^{11,12,13,14,15} predictions. The results are illustrated in Fig. 3 (left, middle), also for the ratio between charge asymmetry ratio

$$A^{+-}(\text{WZ}) = \frac{\sigma_{\text{fid}}(\text{pp} \rightarrow \text{W}^+\text{Z})}{\sigma_{\text{fid}}(\text{pp} \rightarrow \text{W}^-\text{Z})}. \quad (2)$$

The asymmetry ratio is a consequence of the asymmetry in the up and down quark proton PDFs⁴. We therefore use our measurement to compute a Bayesian posterior predictions for the LHC proton PDFs⁴, constraining the uncertainty in their prediction by about 10% with respect to the theoretical value, as illustrated in Fig. 3 (right).

The helicity of the W boson has been measured in several production modes, such as $t\bar{t}$, single top, and W +jets, and by ATLAS in WZ production¹⁰. A measurement in WZ production was difficult so far because of the limited amount of data events available in the WZ signal regions.




Figure 3 – Cross section measurements for WZ production (left) and charge asymmetry (middle). Constraints on the proton PDFs⁴ uncertainty obtained via Bayesian reweighting (right). Figures reproduced from the CMS paper².

With the full Run 2 dataset, we have been able to measure the polarization of both the W and the Z boson¹⁶, by relying on assigning each lepton to its parent boson (which we are able to do with about 95% efficiency), and reconstructing the angle between the parent boson and the resulting lepton, based on templates from simulation. At leading order and assuming no interference, we express the polarization angles as:

$$\frac{1}{\sigma} \frac{\sigma}{\cos \theta^{W\pm}} = \frac{3}{8} \left\{ [1 \mp \cos(\theta^{W\pm})]^2 f_L^W + [1 \pm \cos(\theta^{W\pm})]^2 f_R^W + 2 \sin^2(\theta^{W\pm}) f_0^W \right\} \quad (3)$$

and

$$\frac{1}{\sigma} \frac{\sigma}{\cos \theta^Z} = \frac{3}{8} \left\{ [1 + \cos^2(\theta^Z) - 2c \cos(\theta^Z)] f_L^Z + [1 + \cos^2(\theta^Z) + 2c \cos(\theta^Z)] f_R^Z + 2 \sin^2(\theta^Z) f_0^Z \right\}, \quad (4)$$

as a function of the three polarization fractions in the helicity frame. A corresponding equation governs the polarization angle for the Z boson. A maximum likelihood fit of the polarization angle is performed to determine the three fractions and the overall WZ normalization, with the constraint that the fractions must sum up to unity. Our paper² reports the results in tabular and graphical forms for one-, two-, and three-dimensional fits were the remaining free parameters are set to their SM prediction. The results are generally in agreement with predictions from POWHEG^{11,12,13,14,15} and MATRIX³.

Differential cross sections are measured as a function of several observables: besides those already employed in our previous paper⁷, we also probed the polarization angles of the W and Z bosons, and the jet multiplicity (which is a probe for the validity of the jet simulation). The measurement generally favour the predictions from MATRIX^{3,17}, and are illustrated and tabulated² both in merged form and split by leptonic charge and flavour.

Finally, constraints on anomalous values of the WWZ trilinear gauge coupling are set. WZ production is sensitive to multiple BSM effects as effective low energy theories. We use a generic model with three couplings, that the SM predicts to have the values $g_1^Z = 1$, $k_Z = 1$, $\lambda_Z = 0$. We are less sensitive to the k_Z term, because in WZ production we have no access to the p_T of the W propagator. Deviations from SM values of the couplings are visible at high p_T , and we therefore set constraints on the couplings by performing a maximum likelihood fit of the $M_{\ell\ell p_T^{\text{miss}}}$ distribution: the sensitivity to EFT effects comes from the tails of that distribution. One-, two-, and three dimensional confidence regions (fixing to their SM predictions the parameters that are not determined in each fit) are tabulated and illustrated in the paper². We find to evidence for anomalous values of the couplings. Small correlations between the EFT parameters are inferred from the two-dimensional plots shown in Fig. 4. The paper² also contains the results converted to the Warsaw basis.

In EFTs for WZ production, interference terms between the SM and beyond-SM physics are described in dimension six operators by linear terms that are Λ^{-2} - and Λ^{-4} -suppressed.




Figure 4 – Two-dimensional confidence regions for pairs of couplings in the helicity frame. Correlations between the parameters can be inferred from the elliptical shape and inclination of the regions. Figures reproduced from the CMS paper².




Figure 5 – Example evolution of the confidence regions on one EFT parameter as a function of the cutoff scale for EFT suppression at high energies. Figures reproduced from the CMS paper².

Dimension-eight operators would introduce an additional interference term, and the results would be accurate only up to Λ^{-2} -suppressed contributions. We performed additional maximum likelihood fits by using only the linear terms of the quadratic fitting functions, to check the effect of dropping Λ^{-4} terms from the modelling. This is an important cross check of the structure of EFT theories, and the results are illustrated and tabulated in the paper². Finally, dimension-6 operators lead to nonphysical results characterized by unitarity breaking leading to cross section values of infinity at arbitrarily high energies. We introduce a cutoff scale to suppress EFT at high energies, and show the evolution of the confidence regions for the EFT parameters as a function of the cutoff in Fig. 5.

4 Summary

In this contribution, I have outlined recent precision measurements of the standard model (SM) multiboson production at CMS. A study of diboson production at 5 TeV¹ constitutes an important probe of the SM at a new energy, and the data favour NNLO predictions obtained by MATRIX³. A study of WZ production at 13 TeV² constitutes the most comprehensive study of WZ production to date, containing inclusive and differential cross section measurements, charge asymmetry measurements, constraints on the LHC proton PDFs⁴, and constraints on anomalous values of the WWZ trilinear gauge coupling. No evidence for new physics is found, and all the results favour SM predictions calculated at NNLO using MATRIX^{3,17}.

References

1. The CMS Collaboration, *Phys. Rev. Lett.* **127**, 191801 (2021)
2. The CMS Collaboration, *arXiv* **2110.11231**, (submitted to *JHEP*)
3. M. Grazzini, S. Kallweit, J.M. Lindert, S. Pozzorini, M. Wiesemann, *JHEP* **2020**, 1 (2019)
4. The NNPDF Collaboration *EPJ C* **77**, 663 (2017)

5. The CMS Collaboration, *EPJ C* **81**, 278 (2021)
6. The CMS Collaboration, *arXiv* **2106.14246**, (submitted to *JHEP*)
7. The CMS Collaboration, *JHEP* **04**, 122 (2019)
8. The CMS Collaboration, *EPJ C* **81**, 378 (2021)
9. Carlos Erice Cid, CERN-THESIS-2021-154 , (2021)
10. The ATLAS Collaboration, *EPJ C* **79**, 535 (2019)
11. P. Nason, *JHEP* **11**, 078 (2011)
12. S. Frixione, P. Nason, C. Oleari, *JHEP* **11**, 070 (2007)
13. S. Alioli, P. Nason, R. Rontsch, G. Zanderighi, *JHEP* **06**, 043 (2010)
14. T. Melia, P. Nason, R. Rontsch, G. Zanderighi, *JHEP* **11**, 078 (2011)
15. P. Nason, G. Zanderighi *EPJ C* **74**, 2702 (2014)
16. J. Baglio, L. Duc Ninh, *JHEP* **1904**, 065 (2019)
17. M. Grazzini, S. Kallweit, M. Wiesemann, *EPJ C* **78**, 1 (2017)